

**Purdue University**  
**Purdue e-Pubs**

---

LARS Symposia

Laboratory for Applications of Remote Sensing

---

1-1-1979

# Use of a Standard Deviation Based Texture Channel for Landsat Classification of Forest Strata

Thomas L. Logan

Alan H. Strahler

Curtis E. Woodcock

Follow this and additional works at: [http://docs.lib.purdue.edu/lars\\_symp](http://docs.lib.purdue.edu/lars_symp)

---

Logan, Thomas L.; Strahler, Alan H.; and Woodcock, Curtis E., "Use of a Standard Deviation Based Texture Channel for Landsat Classification of Forest Strata" (1979). *LARS Symposia*. Paper 305.  
[http://docs.lib.purdue.edu/lars\\_symp/305](http://docs.lib.purdue.edu/lars_symp/305)

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

Reprinted from

**Symposium on**

**Machine Processing of**

**Remotely Sensed Data**

**June 27 - 29, 1979**

The Laboratory for Applications of  
Remote Sensing

Purdue University  
West Lafayette  
Indiana 47907 USA

IEEE Catalog No.  
79CH1430-8 MPRSD

Copyright © 1979 IEEE  
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to [pubs-permissions@ieee.org](mailto:pubs-permissions@ieee.org).

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

# USE OF A STANDARD DEVIATION BASED TEXTURE CHANNEL FOR LANDSAT CLASSIFICATION OF FOREST STRATA

THOMAS L. LOGAN  
Informatics, Inc.  
Pasadena, California

ALAN H. STRAHLER  
CURTIS E. WOODCOCK  
Department of Geography,  
University of California,  
Santa Barbara

A classification procedure for identifying volume homogeneous timber strata from Landsat imagery is under examination in the rugged forest lands of northern California's Klamath National Forest. A unique feature of this procedure is the addition of a texture channel created from Landsat Band 5 by calculating the standard deviation of density values within a 3-by-3 pixel moving window. Once calculated, the standard deviation is scaled, associated with the center location of the 3-by-3 window, and output in spatially registered image format. This synthesized texture channel exhibits low values in areas of continuous canopy cover and higher values in areas of discontinuous canopy. Highest values occur at abrupt vegetation boundaries.

When combined with the conventional four Landsat bands in an unsupervised classification, the texture channel promotes discrimination of numerous site-specific classes. These classes are merged as necessary to produce a strata map appropriate for locating the random samples necessary for volume inventory. Use of this Landsat-based procedure should be very effective in reducing the high cost of compiling strata maps from conventional manual procedures involving airphoto interpretation.